



Research

Cite this article: Bedgood SA, Mastroni SE, Bracken MES. 2020 Flexibility of nutritional strategies within a mutualism: food availability affects algal symbiont productivity in two congeneric sea anemone species. *Proc. R. Soc. B* **287**: 20201860.
<https://doi.org/10.1098/rspb.2020.1860>

Received: 2 August 2020

Accepted: 16 November 2020

Subject Category:

Ecology

Subject Areas:

ecology, physiology

Keywords:

Anthopleura, context-dependent, ecophysiology, mutualistic symbiosis, sea anemone, Symbiodiniaceae

Author for correspondence:

Samuel A. Bedgood

e-mail: sbedgood@uci.edu

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.5227819>.

Flexibility of nutritional strategies within a mutualism: food availability affects algal symbiont productivity in two congeneric sea anemone species

Samuel A. Bedgood¹, Sarah E. Mastroni² and Matthew E. S. Bracken¹

¹Department of Ecology and Evolutionary Biology, University of California, 321 Steinhaus Hall, Irvine, CA 92697, USA

²Coastal Science and Policy Program, University of California Santa Cruz, 115 McAllister Way, Santa Cruz, CA 95060, USA

SAB, 0000-0002-2510-0900; MESB, 0000-0002-0068-7485

Mutualistic symbioses are common, especially in nutrient-poor environments where an association between hosts and symbionts can allow the symbiotic partners to persist and collectively out-compete non-symbiotic species. Usually these mutualisms are built on an intimate transfer of energy and nutrients (e.g. carbon and nitrogen) between host and symbiont. However, resource availability is not consistent, and the benefit of the symbiotic association can depend on the availability of resources to mutualists. We manipulated the diets of two temperate sea anemone species in the genus *Anthopleura* in the field and recorded the responses of sea anemones and algal symbionts in the family Symbiodiniaceae to our treatments. Algal symbiont density, symbiont volume and photosynthetic efficiency of symbionts responded to changes in sea anemone diet, but the responses depended on the species of sea anemone. We suggest that temperate sea anemones and their symbionts can respond to changes in anemone diet, modifying the balance between heterotrophy and autotrophy in the symbiosis. Our data support the hypothesis that symbionts are upregulated or downregulated based on food availability, allowing for a flexible nutritional strategy based on external resources.

1. Introduction

In nutrient-poor environments, mutualistic symbioses are common [1–3]. In these symbioses a diverse set of nutrients are exchanged between partners, but the unifying theme is an exchange of carbon and nitrogen. For example, in relatively nutrient-poor environments, partnerships form between legumes and rhizobia [4], fungi and algae (i.e. lichens [5]), and corals and algal endosymbionts [6]. However, these environments are not static, and as resources for hosts and symbionts fluctuate [7,8], the benefit to each partner may change, potentially disrupting the symbiosis. Legumes in nitrogen-enriched soil no longer benefit from their symbiotic rhizobia [8], lichens are impacted by nitrogen deposition [7] and coral–algal symbioses may break down as a result of human-induced nutrient fluctuations [9]. Most previous studies have focused on anthropogenic changes in nutrient availability; we know less about how natural fluctuations in resources affect mutualistic symbioses *in situ*. A species that can obtain external resources when they are plentiful and simultaneously maintain its association with symbionts could employ a flexible nutritional strategy that depends on resource availability.

Scleractinian corals and their algal endosymbionts have been described using an ecophysiological framework based on nutrient and energy exchange since these relationships were first described [10,11]. Studies of coral–algal symbioses have informed our understanding of metabolic exchange between symbiotic

partners including autotrophic products from the algae and heterotrophic nutrients from zooplankton captured by the coral [12–14]. In recent years, a large body of research has focused on the breakdown between corals and their algal symbionts, highlighting the importance of symbionts in coral metabolism [15,16]. However, symbiotic coral species are obligate mutualists (with the exception of *Astrangia poculata*) where symbiont and host derived nutrition are balanced and critical for survival; flexibility between autotrophic and heterotrophic nutritional pathways is limited (but see [16–18]).

Some tropical and temperate sea anemone species are similar to corals in obligately associating with algal endosymbionts [19], but many symbiotic sea anemones, especially temperate species, are facultative mutualists [20]. In contrast to the nutrient-poor environments where corals and some tropical sea anemones live, temperate anemones often benefit from nutrient-rich environments where prey are abundant [19,21], enhancing the potential for nutritional flexibility in these symbioses. Symbiont densities in natural populations can vary substantially, and these densities are affected by light intensity and temperature [22,23] (S.A.B. 2018, unpublished data). At the same time, sea anemones are opportunistic passive suspension feeders that rely on water currents, tides, waves and chance to deliver potential prey, so food availability can be unpredictable and can vary among individuals and across time [24–26]. Whereas several studies have addressed how starvation affects the relationship between anemones and algal symbionts in laboratory manipulations of tropical [27–29] and temperate [30] species, the applicability of these studies to field conditions remains unknown, as little is known about how variation in food availability affects algal symbionts and their contribution to the host sea anemones in the field. If the relationship between the sea anemone and its algal symbionts is driven by the requirements of the anemone host, then symbionts would be downregulated when prey are readily available and upregulated when prey are scarce. Here we investigate if realistic, *in situ* changes in the food available to sea anemone hosts, based on naturally occurring fluctuations observed in previous studies [24] (S.A.B. 2018, unpublished data), affect the abundance, photo-physiology and interactions between algal symbionts and their host sea anemone.

We studied *Anthopleura sola* and *Anthopleura xanthogrammica*, two sea anemone species that host algal symbionts. Both species coexist on California rocky shores [31,32] (S.A.B. 2017, unpublished data), where light is abundant for photosynthesizing symbionts, and food is washed in from adjacent intertidal habitats and the ocean. Both species are similar in size, consume the same prey and use similar habitat in the mid-intertidal zone (S.A.B. 2017, unpublished data; this study).

The algal symbionts within *A. sola* and *A. xanthogrammica* at our study location are in the family Symbiodiniaceae, the same group that includes symbionts in tropical corals [33]. These symbionts are in the genus *Breviolum* (previously Clade B) [34–36] and provide a substantial portion of the anemones' dietary carbon as demonstrated by stable isotope analyses [37,38]. Genetic differences between symbionts in *A. sola* and *A. xanthogrammica* at the same site and tidal height are minimal in this region; genetically identical symbionts are found in both sea anemone species [36]. Therefore, differences in the responses of symbionts are likely to be due to differences between sea anemone species, not differences in symbiont identity.

The growth rate potential of symbiont cells is probably always higher than that of host cells in cnidarian–algal symbioses, so it is crucial that the host has some control of symbiont density [39]. Algal symbionts reproduce asexually within their anemone hosts resulting in higher densities [37] and can vary in volume, probably based on productivity [37,40]. *Anthopleura elegantissima* (a congeneric co-occurring species) can exocytose and ingest algal cells to control their densities [41,42]. There are costs to maintaining high symbiont densities in this species, most notably the production of oxygen radicals (H_2O_2) by photosynthesizing symbionts under intense light that damage host cells [43,44]. While the mechanisms underlying control of symbiont densities in *Anthopleura* spp. are not fully understood, symbiont densities are known to be maintained by nitrogen availability within the host anemone [45,46], by coregulation of host and symbiont cell cycles [47], and by symbiont degradation within the host in tropical cnidarian–algal symbioses [39]. While the algal symbionts may increase their densities by reproducing within the host, the anemone probably has substantial control of symbiont density.

If symbionts function as a partial substitute for captured prey, and there is a cost to the host of maintaining high densities of symbionts within the tissue, then we would expect to observe reduced symbiont abundances when prey are abundant and/or higher abundances when prey are scarce (figure 1). We hypothesize that this symbiotic partnership is nutritionally flexible and therefore predict that realistic changes in host diet will influence three measures of symbiont productivity (figure 1). (i) Symbiont density—which we hypothesize is controlled by the host—will increase when prey are removed and decrease when prey are added. (ii) Individual symbiont cell volume will decrease when prey are removed (i.e. more photosynthetic products are given to the host and less is stored in the symbiont cell) and increase when prey are added (i.e. symbionts store photosynthetic products that are not translocated to the host, increasing cell volume). (iii) Photosynthetic efficiency will be affected by nitrogen availability within the host (i.e. hosts with added prey may translocate more nitrogen to their symbionts). However, we do not predict any change in photosynthetic efficiency when prey are removed, as hosts in nutrient-rich environments are likely to retain nitrogen when prey are scarce.

2. Methods

(a) Site description and experimental treatments

Individuals of both sea anemone species (*A. sola* and *A. xanthogrammica*; $n = 28$ each) were located in the intertidal zone at Kenneth S. Norris Rancho Marino Reserve (35°32'24.32" N, 121° 5'34.12" W). Sea anemones were excluded if their largest closed crown diameter was less than 40 mm because anemones smaller than this had distinctly different diets (i.e. no mussels or sea urchins; S.A.B. 2018, personal observation). We used the length and width of the closed crown to calculate the area (using an ellipse shape) as a measure of anemone size at the beginning and end of the experiment. All sea anemones were located between +0.4 m and +1.1 m above mean lower-low water. Each *A. sola* was paired with a nearby *A. xanthogrammica* within the same habitat. We used a blocked design consisting of 8 sea anemones (4 *A. sola* and 4 *A. xanthogrammica*) in close proximity (e.g. within the same tide pool) that matched all four feeding and species treatments ($n = 7$ blocks).




Figure 2. Boxplots with overlaid data points ($n = 3$ or 4) showing $\delta^{13}\text{C}$ (‰) values for *A. sola*, *A. xanthogrammica* and supplemented prey items. Tissue samples from anemones were separated into anemone and algal symbiont portions before analysis. Comparisons are made among the control, reduction and supplement treatments. Significant main effects and interactions from GLMMs are listed in the upper right-hand corner of each graph with p -values. Asterisks above the x -axis signify significant differences between portions within a treatment from a Tukey *post hoc* analysis. Lower case letters represent significant differences among treatment groups within the algal symbiont portion. (Online version in colour.)

these samples were collected at the end of the experiment. Samples were homogenized as described previously. The homogenate was then centrifuged at 2000g for 5 min to separate the anemone cells from the algal symbiont cells. The top layer of anemone cells was then agitated, and the supernatant with suspended anemone cells was removed. Both the algae portion and anemone portion (supernatant) were re-homogenized and centrifuged 2–3 more times to remove any non-target cells. Both the symbiont and anemone portions were placed on separate microscope slides and dried (60°C for greater than 48 h) before analysis at the UCI Stable Isotope Ratio Mass Spectrometry Facility.

(f) Statistical analyses

We conducted all analyses in R v. 3.6.2 and RStudio 1.2.5003 [52] using the packages lme4 to create general linear mixed models (GLMMs) and emmeans for *post hoc* analyses. We checked the diet composition data for normality with a Shapiro–Wilk test, and then used a paired t -test to compare anemone diets. We used GLMMs paired with ANOVA and Tukey *post hoc* analyses to analyse $\delta^{13}\text{C}$, symbiont density, symbiont cell volume and photosynthetic efficiency. Data from the two anemone species were typically analysed separately. $\delta^{13}\text{C}$ values were analysed using GLMMs with the main effects of treatment and tissue type (anemone and symbiont) and a random effect of anemone. Symbiont density, symbiont cell volume and photosynthetic efficiency were measured over time with two control groups, so we compared treatment groups in pairs through time: control/supplement and probe/reduction. These data were analysed with GLMMs with main effects of treatment and time and a random effect of anemone.

3. Results

(a) Composition of diets

Prey were found in the gastrovascular cavity of *A. xanthogrammica* almost twice as frequently as in *A. sola* (paired t -test:

$t = -3.56$, $p = 0.003$). Prey were found within *A. sola* during $12.92 \pm 2.31\%$ (mean \pm s.e.) of daily checks, while prey were found within *A. xanthogrammica* during $23.47 \pm 3.18\%$ of checks. The greatest proportion of both species' diets (40% of observations) was composed of the California sandcastle worm, *Phragmatopoma californica*. Other prey items included limpets, hermit crabs and sea urchins, but each of these comprised less than 10% of diets. There was no apparent difference in the diet composition of the two anemone species. The frequency of prey was 0.90 ± 0.22 items per week for *A. sola* and 1.64 ± 0.15 items per week for *A. xanthogrammica*. We removed an average of 2–6 items from each anemone in the 'reduction' treatment over the course of the experiment. The diet supplement treatments received an additional prey item daily, which represented a substantial increase from ambient prey capture rates. However, this frequency of food availability is not uncommon during periods of high wave exposure, when all anemones surveyed had at least one prey item on consecutive days.

(b) Stable isotope analysis

Anemone diet affected $\delta^{13}\text{C}$ values (GLMM ANOVA: *A. s.* = *A. sola* treat \times portion, $F = 5.73$, $p = 0.025$; *A. x.* = *A. xanthogrammica* treat, $F = 9.74$, $p = 0.007$), but this result was largely associated with the algal symbiont portion for both species (figure 2). Symbionts from 'supplemented' anemones had $\delta^{13}\text{C}$ signatures that were 2–5‰ lower than the controls (GLMM Tukey HSD: *A. s.*, $t = -4.89$, $p = 0.001$; *A. x.*, $t = -4.1$, $p = 0.004$), but reduction of diet had no effect (*A. s.*, $t = 1.96$, $p = 0.165$; *A. x.*, $t = -0.84$, $p = 0.684$). $\delta^{13}\text{C}$ values did not differ between anemones and their algae within a treatment, except in the supplement treatment where the symbionts had a lower $\delta^{13}\text{C}$ (*A. s.*, $t = 3.0$, $p = 0.015$; *A. x.*, $t = 2.57$, $p = 0.033$).

(c) Symbiont density and chlorophyll *a*

The symbiont density was affected by treatment (GLMM ANOVA: *A. s.*, $F = 5.06$, $p = 0.044$) and the effect of treatment changed over time (*A. x.*, $F = 7.69$, $p = 0.003$), but the effect was observed in different treatment groups in each anemone species. In *A. sola*, supplementing food resulted in decreased symbiont densities after one week of treatment (GLMM Tukey HSD: $t = 2.74$, $p = 0.01$), but symbiont density did not increase when food was reduced ($t = 1.39$, $p = 0.173$). In *A. xanthogrammica*, supplementing food did not affect symbiont density ($t = -0.17$, $p = 0.869$), but reducing food increased symbiont density after one week of treatment ($t = -4.23$, $p < 0.001$). All symbiont density measurements changed over time (figures 3 and 4) due to an increase in symbiont density after one week. Chl *a* per symbiont was not affected by treatment (GLMM ANOVA: *A. s.* reduction, $F = 0.83$, $p = 0.378$; *A. s.* supplement, $F = 0.63$, $p = 0.444$; *A. x.* reduction, $F = 0.17$, $p = 0.684$; *A. x.* supplement, $F = 1.37$, $p = 0.264$), so Chl *a* concentrations tracked symbiont density measurements closely throughout the experiment (figures 3 and 4). However, while there was no effect of supplementation on symbiont density in *A. xanthogrammica*, the Chl *a* concentration in the supplement treatment was lower than the control at week three (figure 4; GLMM Tukey HSD: $t = 2.55$, $p = 0.017$). Anemone growth (final size – initial size/initial size) was not different among treatment groups at the final time point (ANOVA: *A. s.* reduction, $F = 0.28$, $p = 0.607$; *A. s.* supplement, $F = 0.29$, $p = 0.603$; *A. x.* reduction, $F = 1.97$, $p = 0.186$; *A. x.* supplement, $F = 2.95$, $p = 0.112$), so anemone growth did not affect symbiont density measurements asymmetrically among groups.

(d) Symbiont cell volume

Both sea anemone species had larger symbionts in the supplement treatment (GLMM Tukey HSD: *A. s.*, $t = -4.69$, $p < 0.001$; *A. x.*, $t = -2.26$, $p = 0.033$; figures 3 and 4) and symbionts were marginally smaller in *A. xanthogrammica* where food was reduced ($t = 2.05$, $p = 0.051$). There was a main effect of time in both species and treatment comparisons where symbiont volume generally decreased over the course of the experiment (*A. s.* reduction, $F = 25.1$, $p < 0.001$; *A. s.* supplement, $F = 17.2$, $p < 0.001$; *A. x.* reduction, $F = 11.5$, $p < 0.001$; *A. x.* supplement, $F = 4.20$, $p = 0.028$).

(e) Photosynthetic efficiency

The photosynthetic efficiency of algal symbionts was higher in *A. sola* than in *A. xanthogrammica* at the start of the experiment (paired *t*-test: $t = 5.72$, $p < 0.001$). This difference persisted throughout the experiment, except when food was supplemented. Then, photosynthetic efficiency in *A. xanthogrammica* increased from 0.56 ± 0.05 (mean \pm s.e.) to 0.71 ± 0.01 F_v/F_m (GLMM Tukey HSD: $t = -2.91$, $p = 0.006$) and did not differ from the mean photosynthetic efficiency of the control treatment *A. sola* symbionts (0.67 ± 0.01 F_v/F_m) by the end of the experiment (paired *t*-test: $t = -1.35$, $p = 0.225$). Photosynthetic efficiency generally increased through time for all groups (figures 3 and 4; GLMM ANOVA: *A. s.* supplement, $F = 5.61$, $p = 0.01$; *A. x.* reduction, $F = 5.41$, $p = 0.012$; *A. x.* supplement, $F = 8.07$, $p = 0.002$) except the *A. sola* reduction pairing ($F = 1.74$, $p = 0.198$).

4. Discussion

Algal symbionts within two species of sea anemone responded to changes in anemone diet, but the responses differed between the anemone species and changed over the course of the experiment. Our framework for dietary carbon-source switching (figure 1) was supported by our results, but support for our predictions depended on the anemone species. Symbionts within *A. sola* responded to diet supplementation, and symbionts within *A. xanthogrammica* responded to both reduction and supplementation. This may be associated with the fact that *A. xanthogrammica* captured twice as many prey items as *A. sola*, so the reduction treatment had a larger impact on *A. xanthogrammica* than on *A. sola*. Supplementation affected both species, resulting in reduced $\delta^{13}\text{C}$ values in symbionts. Furthermore, $\delta^{13}\text{C}$ did not differ between anemones and their symbionts, except where food was added. Lower $\delta^{13}\text{C}$ values have previously been associated with an increase in heterotrophy in corals [53] and in *Anthopleura* anemones [37,38]. A lower $\delta^{13}\text{C}$ signature (supplement treatment) occurs when algae selectively incorporate the lighter carbon isotope (^{12}C) over the heavier isotope (^{13}C). Highly productive algal symbionts at high densities cannot choose the lighter carbon isotope because CO_2 is limited within the host tissue, resulting in a heavier carbon isotope signature (reduction and control treatments) [54].

Symbiont densities were affected by host dietary changes, but underlying mechanisms are not well understood. It is likely that the sea anemone host benefits from a reduction in symbiont density when they are unnecessary (supplement treatment) as they can cause damage to tissue via oxygen radicals [43,44]. The host would also benefit from an increase in symbiont density or chlorophyll when heterotrophic diet decreases (reduction treatment) to compensate for lost dietary carbon as an increase in either would allow for increased translocation of photosynthetic products from the symbionts to the host (figure 1).

The anemone–algae holobiont responded to supplementation of the host diet largely by decreasing symbiont density and/or chlorophyll while increasing symbiont cell volume. This could have resulted from egestion of symbionts or by the slowing of symbiont reproduction within the host. The remaining symbionts may have been larger because they were able to store resources rather than translocate them to the host or because they did not asexually reproduce. More research is needed to fully understand the mechanism(s) driving symbiont volume changes in these anemones. Regardless of the mechanisms, those anemones that received more external resources (prey) had lower autotrophic potential (fewer symbionts and/or lower chlorophyll). However, symbionts within *A. xanthogrammica* may have compensated for the decrease in chlorophyll by increasing photosynthetic efficiency.

Reduction of host diet had an effect on *A. xanthogrammica* and its symbionts but not on *A. sola*. Symbiont density increased and symbiont volume decreased when food was reduced in *A. xanthogrammica*, suggesting that the anemone host maintained a higher symbiont density to compensate for the loss of dietary carbon by either retaining symbionts that would otherwise be egested or by increasing the reproduction of symbionts. *A. xanthogrammica* anemones that received fewer external resources had higher autotrophic potential (symbiont density and chlorophyll), but the effect was short-lived and disappeared after three weeks of treatment.




Figure 3. Boxplots showing symbiont density, chlorophyll concentration, symbiont cell volume and photosynthetic efficiency of symbionts within *A. sola* throughout the experiment. Comparisons are made between the supplement and reduction treatments and their respective control treatments. The vertical dashed line represents the start of treatments. Significant main effects and interactions from GLMMs are listed in the upper right-hand corner of each graph with p -values. Asterisks above the x -axis signify significant differences between controls and treatments at a given time point from a Tukey *post hoc* analysis. $n = 7$ for each treatment at each time point. (Online version in colour.)

Our results suggest there is a trade-off between sources of nutrition—external and symbiont-mediated—in this mutualism. Similar previous work that involved starving sea

anemones under laboratory conditions provided conflicting perspectives on the effect of host diet on symbiont density [29,55,56], but we show here that realistic, *in situ*




Figure 4. Boxplots showing symbiont density, chlorophyll concentration, symbiont cell volume and photosynthetic efficiency of symbionts within *A. xanthogrammica* throughout the experiment. Comparisons are made between the supplement and reduction treatments and their respective control treatments. The vertical dashed line represents the start of treatments. Significant main effects and interactions from GLMMs are listed in the upper right-hand corner of each graph with p -values. Asterisks above the x -axis signify significant differences between controls and treatments at a given time point from a Tukey *post hoc* analysis. $n = 7$ for each treatment at each time point. (Online version in colour.)

changes in sea anemone diet reveal ecologically relevant trade-offs in symbiont–host nutrition that were previously unexplored.

Not all algae-hosting cnidarians can switch carbon sources. Tropical corals tend to lose symbionts when starved [57,58], suggesting that symbionts do not serve as a comparable

nutritional pathway in the absence of heterotrophy (but see [59]). This is probably because most tropical corals are obligate mutualists, whereas *Anthopleura* anemones are facultative. A better comparison may be to a freshwater hydra where algal symbiont density decreases immediately after predatory feeding [60] and increases with starvation [61].

Analogous partner interactions exist in terrestrial mutualisms where legumes host fewer rhizobium (via nodules) when external sources of nitrogen are available in the soil [62,63] and the benefit and cost of arbuscular mycorrhizal fungi to plants is dependent on environmental resources [64]. Holobionts with flexible nutritional strategies—like the ones we describe here—may be able to withstand periods of resource limitation, allowing species to persist in an otherwise inhospitable environment. Interactions between hosts and symbionts are dependent on external resource availability in normally nutrient-poor environments. Some mutualisms may break down as a result of perturbations [8,65], but others are flexible, requiring more from symbionts when nutrients are scarce or less from them when nutrients are abundant [60,61,66]. Future research on flexible mutualisms should focus on how realistic fluctuations of external resources affect the production and storage of resources by symbiotic partners.

Our results suggest that even modest changes in resource availability have the potential to alter the interaction between partners in a mutualistic symbiosis, but those changes are species-specific even in congeneric species sharing the same symbiont. We found evidence for a trade-off between autotrophic and heterotrophic nutritional pathways within an algal-symbiont-hosting sea anemone, but these pathways are not equal. We propose that autotrophy allows for persistence, but growth probably requires heterotrophy as evidenced in this and other studies on cnidarians [61,67].

Anemone hosts and algal symbionts respond to changes in heterotrophic diet by altering their interactions with each other, compensating for externally derived nutrition. The potential for flexible nutritional strategies in other mutualistic symbioses is largely unexplored, especially in systems where environmental resources are naturally stochastic.

Ethics. All applicable international, national and/or institutional guidelines for the care and use of animals were followed. This work was conducted under a California Department of Fish and Wildlife Scientific Collecting Permit to S.A.B. (ID number SC-13728).

Data accessibility. Dataset files and R code for analyses and graphs are available from the Dryad Digital Repository: <https://doi.org/10.7280/D1GD62> [68].

Authors' contributions. S.A.B. conceived of the study, designed the study, carried out the study, analysed the data and drafted the manuscript; S.E.M. carried out the study and made contributions to the manuscript; M.E.S.B. advised on experimental design, assisted in analysing the data and added major intellectual contributions to the manuscript. All authors gave final approval for publication and agree to be held accountable for the work performed therein.

Competing interests. We declare we have no competing interests.

Funding. This research was made possible through funding provided by the UC Irvine Oceans, Changing Environments, Arts, and Nearshore Societies (UCI OCEANS) Fellowship (to S.A.B.) and the National Science Foundation (Graduate Research Fellowship to S.A.B. and OCE-1736891 to M.E.S.B. and A. Martiny). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. All applicable state and institutional guidelines for the care and use of animals were followed.

Acknowledgements. We thank the late Don Canestro for the support he provided to us at Kenneth S. Norris Rancho Marino Reserve; this research would not have been possible without his assistance. We thank Joshua Goldberg and Nyssa Silbiger for their advice on analysis of our data, and Samantha Levell for her comments on early drafts of this manuscript. We also thank three anonymous reviewers who greatly improved the final version of this text.

References

- Muscatine L, Porter JW. 1977 Reef corals: mutualistic symbioses adapted to nutrient-poor environments. *Bioscience* **27**, 454–460. (doi:10.2307/1297526)
- Boucher DH, James S, Keeler KH. 1982 The ecology of mutualism. *Annu. Rev. Ecol. Syst.* **13**, 315–347. (doi:10.1146/annurev.es.13.110182.001531)
- Odum EP, Biever LJ. 1984 Resource quality, mutualism, and energy partitioning in food chains. *Am. Nat.* **124**, 360–376. (doi:10.1086/284279)
- Masson-Boivin C, Sachs JL. 2018 Symbiotic nitrogen fixation by rhizobia—the roots of a success story. *Curr. Opin. Plant Biol.* **44**, 7–15. (doi:10.1016/j.pbi.2017.12.001)
- Nash TH. 1996 *Lichen biology*. Cambridge, UK: Cambridge University Press.
- Yellowlees D, Rees TAV, Leggat W. 2008 Metabolic interactions between algal symbionts and invertebrate hosts. *Plant Cell Environ.* **31**, 679–694. (doi:10.1111/j.1365-3040.2008.01802.x)
- van Herk CM, Mathijssen-Spiekman EAM, de Zwart D. 2003 Long distance nitrogen air pollution effects on lichens in Europe. *Lichenologist* **35**, 347–359. (doi:10.1016/S0024-2829(03)00036-7)
- Regus JU, Wendlandt CE, Bantay RM, Gano-Cohen KA, Gleason NJ, Hollowell AC, O'Neill MR, Shahin KK, Sachs JL. 2017 Nitrogen deposition decreases the benefits of symbiosis in a native legume. *Plant Soil* **414**, 159–170. (doi:10.1007/s11104-016-3114-8)
- Muller-Parker G, D'Elia CF, Cook CB. 2015 Interactions between corals and their symbiotic algae. In *Coral reefs in the Anthropocene* (ed. C Birkeland), pp. 99–116. Dordrecht, The Netherlands: Springer.
- Yonge CM. 1931 The significance of the relationship between corals and zooxanthellae. *Nature* **128**, 309–311. (doi:10.1038/128309a0)
- Johannes RE, Coles SL, Kuenzel NT. 1970 The role of zooplankton in the nutrition of some scleractinian corals. *Limnol. Oceanogr.* **15**, 579–586. (doi:10.4319/lo.1970.15.4.0579)
- Davies PS. 1984 The role of zooxanthellae in the nutritional energy requirements of *Pocillopora eydouxi*. *Coral Reefs* **2**, 181–186.
- Davies PS. 1991 Effect of daylight variations on the energy budgets of shallow-water corals. *Mar. Biol.* **108**, 137–144. (doi:10.1007/BF01313481)
- Houlbrèque F, Ferrier-Pagès C. 2009 Heterotrophy in tropical scleractinian corals. *Biol. Rev.* **84**, 1–17. (doi:10.1111/j.1469-185X.2008.00058.x)
- Matthews JL, Crowder CM, Oakley CA, Lutz A, Roessner U, Meyer E, Grossman AR, Weis VM, Davy SK. 2017 Optimal nutrient exchange and immune responses operate in partner specificity in the cnidarian-dinoflagellate symbiosis. *Proc. Natl Acad. Sci. USA* **114**, 13 194–13 199. (doi:10.1073/pnas.1710733114)
- Morris LA, Voolstra CR, Quigley KM, Bourne DG, Bay LK. 2019 Nutrient availability and metabolism affect the stability of coral–Symbiodiniaceae symbioses. *Trends Microbiol.* **27**, 678–689. (doi:10.1016/j.tim.2019.03.004)
- Szmant-Froelich A, Pilson MEQ. 1980 The effects of feeding frequency and symbiosis with zooxanthellae on the biochemical composition of *Astrangia Danae* Milne Edwards & Haime 1849. *J. Exp. Mar. Biol. Ecol.* **48**, 85–97. (doi:10.1016/0022-0981(80)90009-X)
- Dimond J, Carrington E. 2007 Temporal variation in the symbiosis and growth of the temperate scleractinian coral *Astrangia poculata*. *Mar. Ecol. Prog. Ser.* **348**, 161–172. (doi:10.3354/meps07050)

19. Muller-Parker G, Davy SK. 2001 Temperate and tropical algal-sea anemone symbioses. *Invertebr. Biol.* **120**, 104–123.
20. Hiebert TC, Bingham BL. 2012 The effects of symbiotic state on heterotrophic feeding in the temperate sea anemone *Anthopleura elegantissima*. *Mar. Biol.* **159**, 939–950. (doi:10.1007/s00227-011-1871-8)
21. Davy SK, Lucas IAN, Turner JR. 1996 Carbon budgets in temperate anthozoan-dinoflagellate symbioses. *Mar. Biol.* **126**, 773–783. (doi:10.1007/BF00351344)
22. Saunders BK, Muller-Parker G. 1997 The effects of temperature and light on two algal populations in the temperate sea anemone *Anthopleura elegantissima* (Brandt, 1835). *J. Exp. Mar. Biol. Ecol.* **211**, 213–224. (doi:10.1016/S0022-0981(96)02723-2)
23. Secord D, Muller-Parker G. 2005 Symbiont distribution along a light gradient within an intertidal cave. *Limnol. Oceanogr.* **50**, 272–278. (doi:10.4319/lo.2005.50.1.0272)
24. Sebens KP. 1981 Recruitment in a sea anemone population: juvenile substrate becomes adult prey. *Science* **213**, 785–787. (doi:10.1126/science.213.4509.785)
25. Shick JM. 2012 *A functional biology of sea anemones*. New York, NY: Springer Science & Business Media.
26. Wells CD. 2019 Biology and ecology of hexacorallians in the San Juan archipelago. Thesis. See <https://digital.lib.washington.edu/443/researchworks/handle/1773/44063>.
27. Muller-Parker G, Lee KW, Cook CB. 1996 Changes in the ultrastructure of symbiotic Zooxanthellae (*Symbiodinium* sp., Dinophyceae) in fed and starved sea anemones maintained under high and low light. *J. Phycol.* **32**, 987–994. (doi:10.1111/j.0022-3646.1996.00987.x)
28. Cook CB, Muller-Parker G, D'Elia CF. 1992 Ammonium enhancement of dark carbon fixation and nitrogen limitation in symbiotic zooxanthellae: effects of feeding and starvation of the sea anemone *Aiptasia pallida*. *Limnol. Oceanogr.* **37**, 131–139. (doi:10.4319/lo.1992.37.1.0131)
29. Muller-Parker G. 1985 Effect of feeding regime and irradiance on the photophysiology of the symbiotic sea anemone *Aiptasia pulchella*. *Mar. Biol.* **90**, 65–74. (doi:10.1007/BF00428216)
30. Fitt WK, Pardy RL. 1981 Effects of starvation, and light and dark on the energy metabolism of symbiotic and aposymbiotic sea anemones, *Anthopleura elegantissima*. *Mar. Biol.* **61**, 199–205. (doi:10.1007/BF00386660)
31. Hand C. 1955 The sea anemones of central California. Part II. The endomyarian and mesomyarian anemones. *Wasmann J. Biol.* **13**, 37–100.
32. Francis L. 1973 Intraspecific aggression and its effect on the distribution of *Anthopleura elegantissima* and some related sea anemones. *Biol. Bull.* **144**, 73–92. (doi:10.2307/1540148)
33. LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR, Santos SR. 2018 Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Curr. Biol.* **28**, 2570–2580.e6. (doi:10.1016/j.cub.2018.07.008)
34. LaJeunesse TC, Trench RK. 2000 Biogeography of two species of *Symbiodinium* (Freudenthal) inhabiting the intertidal sea anemone *Anthopleura elegantissima* (Brandt). *Biol. Bull.* **199**, 126–134. (doi:10.2307/1542872)
35. Sanders JG, Palumbi SR. 2011 Populations of *Symbiodinium muscatinei* show strong biogeographic structuring in the intertidal anemone *Anthopleura elegantissima*. *Biol. Bull.* **220**, 199–208. (doi:10.1086/BBLv220n3p199)
36. Cornwell BH. 2018 Environmental variation across nested spatial scales differentially shapes patterns of host and symbiont population genetic structure in a marine symbiotic relationship. PhD thesis, University of California, Davis. See <https://search.proquest.com/dissertations/docview/2088559165/10E76F14556D4C3DPQ/1>.
37. Bergschneider H, Muller-parker G. 2008 Nutritional role of two algal symbionts in the temperate sea anemone *Anthopleura elegantissima* (Brandt). *Biol. Bull.* **215**, 73–88. (doi:10.2307/25470685)
38. Levine M, Muller-Parker G. 2012 Distribution patterns and nutritional contributions of algal symbionts in the sea anemone *Anthopleura xanthogrammica*. *Mar. Ecol. Prog. Ser.* **453**, 79–94. (doi:10.3354/meps09602)
39. Davy SK, Allemand D, Weis VM. 2012 Cell biology of cnidarian-dinoflagellate symbiosis. *Microbiol. Mol. Biol. Rev.* **76**, 229–261. (doi:10.1128/MMBR.05014-11)
40. McBride BB, Muller-Parker G, Jakobsen HH. 2009 Low thermal limit of growth rate of *Symbiodinium californium* (Dinophyta) in culture may restrict the symbiont to southern populations of its host anemones (*Anthopleura* spp.; Anthozoa, Cnidaria). *J. Phycol.* **45**, 855–863. (doi:10.1111/j.1529-8817.2009.00716.x)
41. McCloskey LR, Cove TG, Verde EA. 1996 Symbiont expulsion from the anemone *Anthopleura elegantissima* (Brandt) (Cnidaria; Anthozoa). *J. Exp. Mar. Biol. Ecol.* **195**, 173–186. (doi:10.1016/0022-0981(95)00079-8)
42. Muller-Parker G, Pierce-Cravens J, Bingham BL. 2007 Broad thermal tolerance of the symbiotic dinoflagellate *Symbiodinium muscatinei* (Dinophyta) in the sea anemone *Anthopleura elegantissima* (Cnidaria) from northern latitudes. *J. Phycol.* **43**, 25–31. (doi:10.1111/j.1529-8817.2006.00302.x)
43. Dykens J, Shick J, Benoit C, Buettner G, Winston G. 1992 Oxygen radical production in the sea anemone *Anthopleura elegantissima* and its endosymbiotic algae. *J. Exp. Biol.* **168**, 219–241.
44. Dimond JL, Orechovesky S, Oppenheimer J, Rodríguez-Ramos J, Bingham BL. 2017 Photophysiology and hydrogen peroxide generation of the dinoflagellate and chlorophyte symbionts of the sea anemone *Anthopleura elegantissima*. *J. Exp. Mar. Biol. Ecol.* **489**, 43–47. (doi:10.1016/j.jembe.2017.01.008)
45. Smith GJ, Muscatine L. 1999 Cell cycle of symbiotic dinoflagellates: variation in G1 phase-duration with anemone nutritional status and macronutrient supply in the *Aiptasia pulchella*–*Symbiodinium pulchrorum* symbiosis. *Mar. Biol.* **134**, 405–418. (doi:10.1007/s002270050557)
46. Xiang T, Lehnert E, Jinkerson RE, Clowez S, Kim RG, DeNofrio JC, Pringle JR, Grossman AR. 2020 Symbiont population control by host-symbiont metabolic interaction in Symbiodiniaceae–cnidarian associations. *Nat. Commun.* **11**, 108. (doi:10.1038/s41467-019-13963-z)
47. Tivey TR, Parkinson JE, Weis VM. 2020 Host and symbiont cell cycle coordination is mediated by symbiotic state, nutrition, and partner identity in a model cnidarian-dinoflagellate symbiosis. *mBio* **11**, 1–17. (doi:10.1128/mBio.02626-19)
48. Schindelin J *et al.* 2012 Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682. (doi:10.1038/nmeth.2019)
49. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. 1951 Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265–275.
50. Hillebrand H, Dürsel C-D, Kirschtel D, Pollinger U, Zohary T. 1999 Biovolume calculation for pelagic and benthic microalgae. *J. Phycol.* **35**, 403–424. (doi:10.1046/j.1529-8817.1999.3520403.x)
51. Dykens JA, Shick JM. 1984 Photobiology of the symbiotic sea anemone, *Anthopleura elegantissima*: defenses against photodynamic effects, and seasonal photoacclimatization. *Biol. Bull.* **167**, 683–697. (doi:10.2307/1541419)
52. R Core Team. 2019 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. See <https://www.R-project.org/>.
53. Alamaru A, Loya Y, Brokovich E, Yam R, Shemesh A. 2009 Carbon and nitrogen utilization in two species of Red Sea corals along a depth gradient: insights from stable isotope analysis of total organic material and lipids. *Geochim. Cosmochim. Acta* **73**, 5333–5342. (doi:10.1016/j.gca.2009.06.018)
54. Muscatine L, Porter JW, Kaplan IR. 1989 Resource partitioning by reef corals as determined from stable isotope composition. *Mar. Biol.* **100**, 185–193. (doi:10.1007/BF00391957)
55. Clayton WS, Lasker HR. 1984 Host feeding regime and zooxanthellal photosynthesis in the anemone, *Aiptasia pallida* (Verrill). *Biol. Bull.* **167**, 590–600. (doi:10.2307/1541412)
56. Cook CB, D'Elia CF, Muller-Parker G. 1988 Host feeding and nutrient sufficiency for zooxanthellae in the sea anemone *Aiptasia pallida*. *Mar. Biol.* **98**, 253–262. (doi:10.1007/BF00391203)
57. Titlyanov EA, Titlyanova TV, Yamazato K, van Woesik R. 2001 Photo-acclimation of the hermatypic coral *Stylophora pistillata* while subjected to either starvation or food provisioning. *J. Exp. Mar. Biol. Ecol.* **257**, 163–181. (doi:10.1016/S0022-0981(00)00308-7)
58. Towle EK, Enochs IC, Langdon C. 2015 Threatened Caribbean coral is able to mitigate the adverse effects of ocean acidification on calcification by increasing feeding rate. *PLoS ONE* **10**, e0123394. (doi:10.1371/journal.pone.0123394)

59. Grotto AG, Rodrigues LJ, Palardy JE. 2006 Heterotrophic plasticity and resilience in bleached corals. *Nature* **440**, 1186–1189. (doi:10.1038/nature04565)
60. Fishman Y, Zlotkin E, Sher D. 2008 Expulsion of symbiotic algae during feeding by the green hydra—a mechanism for regulating symbiont density? *PLoS ONE* **3**, e2603. (doi:10.1371/journal.pone.0002603)
61. Douglas A, Smith DC. 1984 The green hydra symbiosis. VIII. Mechanisms in symbiont regulation. *Proc. R. Soc. Lond. B* **221**, 291–319. (doi:10.1098/rspb.1984.0035)
62. Arrese-Igor C, Minchin FR, Gordon AJ, Nath AK. 1997 Possible causes of the physiological decline in soybean nitrogen fixation in the presence of nitrate. *J. Exp. Bot.* **48**, 905–913. (doi:10.1093/jxb/48.4.905)
63. Voisin A-S, Salon C, Munier-Jolain NG, Ney B. 2002 Effect of mineral nitrogen on nitrogen nutrition and biomass partitioning between the shoot and roots of pea (*Pisum sativum* L.). *Plant Soil* **242**, 251–262. (doi:10.1023/A:1016214223900)
64. Johnson NC, Graham JH, Smith FA. 1997 Functioning of mycorrhizal associations along the mutualism–parasitism continuum. *New Phytol.* **135**, 575–585. (doi:10.1046/j.1469-8137.1997.00729.x)
65. Wiedenmann J, D'Angelo C, Smith EG, Hunt AN, Legiret F-E, Postle AD, Achterberg EP. 2013 Nutrient enrichment can increase the susceptibility of reef corals to bleaching. *Nat. Clim. Change* **3**, 160–164. (doi:10.1038/nclimate1661)
66. Mariotte P, Canarini A, Dijkstra FA. 2017 Stoichiometric N:P flexibility and mycorrhizal symbiosis favour plant resistance against drought. *J. Ecol.* **105**, 958–967. (doi:10.1111/1365-2745.12731)
67. Bedgood SA, Bracken MES, Ryan WH, Levell ST, Wulff J. 2020 Nutritional drivers of adult locomotion and asexual reproduction in a symbiont-hosting sea anemone *Exaiptasia diaphana*. *Mar. Biol.* **167**, 39. (doi:10.1007/s00227-020-3649-3)
68. Bedgood SA, Mastroni SE, Bracken MES. 2020 Data from: Flexibility of nutritional strategies within a mutualism: food availability affects algal symbiont productivity in two congeneric sea anemone species. Dryad Digital Repository. (doi:10.7280/D1GD62)